

Comprehensive Reviews In Food Science and Food Safety

Flavor Lexicons

M.A. Drake and G.V. Civille

ABSTRACT: Flavor lexicons are a widely used tool for documenting and describing sensory perception of a selected food. Development of a representative flavor lexicon requires several steps, including appropriate product frame of reference collection, language generation, and designation of definitions and references before a final descriptor list can be determined. Once developed, flavor lexicons can be used to record and define product flavor, compare products, and determine storage stability, as well as interface with consumer liking and acceptability and chemical flavor data.

Lexicon characteristics

A flavor lexicon is simply a set of words to describe the flavor of a product or commodity. A lexicon is then applied or practiced using descriptive sensory analysis techniques. A lexicon, like a specific technical dictionary, provides a source list to describe a category of products, such as commodities (cheese, beef, fish) or finished products (bread, orange juice, coffee, chocolate). Although the descriptive panel generates its own list to describe the product array under study, a lexicon provides a source of possible terms with references and definitions for clarification (see ASTM lexicon comments).

As a broad example, the ASTM Lexicon (Civille and Lyon 1996) provides hundreds of words that might be selected by a descriptive panel to describe any number of product categories. The clear admonition for use of the ASTM Lexicon and/or for generation of any lexicon is to:

- (1) collect a product frame of reference
- (2) generate the terms
- (3) review references and examples
- (4) develop a final descriptor list

When a panel begins work in a product category, the 1st source of terminology is the wide sample set or frame of reference. The panel needs to evaluate several (25 to 100) products in the category that cover the widest array of flavor characteristics. The Aromatics, Tastes, and Feeling Factors that the samples manifest are the basis for the lexicon. After some initial attempts at organizing the terms generated by all of the panelists, the panel and panel leader organize references to provide clarification of the grouped terms. Often many terms relate to original ingredients and the processes they are subjected to in the creation of the category. Panelists need to experience these core ingredients (1) with and without process treatments and (2) in different ingredient combinations. After references, the panel goes back to the original list of terms to set the lexicon: merging like terms, eliminating redundant terms, and organizing the list in the order that the attributes appear in most products. This review will include lexicon history, methods, and applications.

Descriptive Analysis Techniques

The 1st published descriptive sensory technique is the Flavor Profile Method (FPM) developed in the 1950s by Arthur D. Little, Inc. (Pangborn 1989; Lawless and Heymann 1998; Meilgaard and others 1999). More recent applications emerged in the 1970s with Quantitative Descriptive Analysis (QDA) and the Spectrum™ method of descriptive analysis. The latter 2 techniques differ markedly from FPM in that they are designed to take measurements from individual panelists and then generate a panel average, rather than generation of a group consensus profile as with FPM (Piggott and others 1998). A detailed overview of all 3 techniques can be found in the Manual on Descriptive Analysis Testing for Sensory Evaluation (1992). Panelist selection is a critical aspect of descriptive analysis (Zook and Wessman 1977; Lawless and Heymann 1998; Meilgaard and others 1999). A good prospective flavor panelist should be able to describe sensations and should be able to discriminate between taste and aroma sensations. Typical screening tests include taste and aroma acuity and discriminatory ability. Good health, enthusiasm, and usage of the general product category are also characteristics of good prospective sensory panelists. All 3 techniques involve screening panelists for sensitivity to flavor and aroma and discriminatory ability followed by extensive training to generate a group of individuals who can function analogously to an instrument to evaluate flavor of products.

In FPM, generally fewer panelists are used, with a minimum of 4 (Keane 1992). The panel leader is a member of the panel and works with the group to generate the language and the method for sample presentation and evaluation. Panelists are screened for sensory acuity, as well as for interest in sensory panel work. Following extensive training (approximately 60 h) panelists work as a group to generate a consensus profile of the sensory properties of the product and the intensity of each. In addition, overall amplitude of the product, as well as a measure of balance and blend are evaluated. The scale used for character (descriptor) notes is a 4 point scale (0-not present, 1-slight, 2-moderate, 3-strong) with half point designations in between, for a total of 7 points of discrimination. Attributes are evaluated in order of appearance. Aftertaste

may also be evaluated. The group consensus process was not designed for statistical analysis, although the scale may be modified to allow statistical analysis.

In QDA, the panel leader is a sensory professional, rather than a member of the panel. During training, the panel (ideally 8 to 12 prescreened individuals) generates the language to describe the product (Stone 1992). The panel leader does not participate in the discussion to generate the flavor attributes, but facilitates the process. The panel also determines the order of appearance of the attributes and, thus, order of evaluation for each descriptor. Definitions for each descriptor are generated and additional products, which may further clarify terms, may also be used. The panel also practices rating products to become familiar with the process and to gain confidence in their abilities. Data is collected on scorecards using line scales anchored on each end. Panelists mark the scale using a dash. The marks are converted to numerical scores by measuring the responses on the scale with a ruler, digitizer, or computerized system. The data can be analyzed statistically and is traditionally graphically represented using a webplot.

In the Spectrum™ method, a universal intensity scale is used (Munoz and Civille 1992; Meilgaard and others 1999). Panelists are screened for cognitive, descriptive and sensory discrimination ability, interest, and availability. The panel size is similar to QDA. By this method, panelists score intensities in the same manner across all attributes. The scales are anchored on either end and can be 15-cm line scales or, more commonly, 0 to 15 numerical scales with tenth subdivisions between, yielding 150 points. The scales and their intensities have already been established, and during training, panelists are given established examples to familiarize them with the use of the scale. Although the scale anchors end at 15, panelists can assign higher numbers if necessary. The panel leader, who should have previous experience with the Spectrum™ method, takes an active role in panel training, including scale usage and language development and application. The panel, with assistance of the panel leader, identifies the sensory language for the product, their order of appearance, and definitions for each term. While a universal intensity scale is standard for the Spectrum™ method, product specific scaling can also be applied. As with QDA, data is readily analyzed by statistical techniques. Results are normally graphically presented using histograms.

Development of Flavor Lexicons

Characteristics of flavor lexicons have been previously discussed by Civille and Lawless (1986) and by Lawless and Heymann (1998). One key characteristic of a good flavor lexicon is that it be discriminating and descriptive. Ideally, for a descriptive language to be discriminating (able to differentiate the products for which it was developed), the language should be developed from a broad representative sample set that exhibits all of the potential variability within the product (Meilgaard and others 1999). Defining what constitutes representative is often a challenge, particularly with foods or commodities which are diverse in flavor, source and treatment. Standard descriptive languages have been developed for commodities using representative sample sets (Jan-to and others 1998; Johnsen and Civille 1986; Johnsen 1986; Johnsen and others 1987; Heisserer and Chambers 1993). In an extreme example, Drake and others (2001b) collected 220 samples of Cheddar cheese varying in age, milk heat treatment, and geographical origin to identify a descriptive language for Cheddar cheese. The sample set was screened to 70 cheeses prior to language generation.

Another key aspect of a flavor lexicon is for the terms to be nonredundant. That is, that multiple terms not be used to describe the same flavor or, alternatively, 1 term representing or overlap-

ping with several other flavors. Redundancy may not be resolved during panel training. Panelists may insist that particular terms are distinct when they are not (Lawless and Heymann 1998). Statistical analysis of results from a representative selection of products is often required to clarify attribute relationships. Drake and others (2001b) reported that the use of the term “aged” in a Cheddar cheese flavor language was in fact a meta term that was comprised of 3 flavor terms and 1 basic taste.

Flavor lexicon terms must also be descriptive with precise and clear meaning for consistent panel performance and for a platform from which research results can be repeated or compared in the future. Panelists may prefer to link descriptors together (for example burnt/charcoal, cooked milk/butterscotch) to assist with group conceptualization of particular attribute. For example, in the USDA Peanut Lexicon (Johnsen 1986), for the terms of woody, hulls, and skins, which represent the more base heavy characteristics inherent under the roast peanut, sweet aromatics, dark roast, and raw beany flavors are combined into 1 term: “woody/hulls/skins.” The use of descriptor definitions and references maximizes language clarity (and minimizes confusion). References may be food, chemicals or other substances. Multiple references are recommended, as different panelists identify better with certain references (Ishii and O’Mahoney 1987, 1991). More importantly, multiple references communicate a “concept,” derived from several references. Single references manifest 1 set of characteristics, so the panel may not be able to relate to the attribute, as seen in some of the products in the category. Definitions are quite useful but may not always provide a frame of reference for all panelists or accurately or fully describe the concept. For instance, the descriptor rancid/free fatty acid is defined as the aromatics associated with short chain free fatty acids. For many panelists unfamiliar with the word or concept, the definition does not necessarily provide a frame of reference. However, when provided with butyric acid (chemical reference) or feta cheese (food reference), a concept and common point of reference becomes readily grasped by all panelists. Meilgaard and others (1982) published a list of chemical references for the beer flavor lexicon. Food or chemical references were also published for 3 of the published cheese lexicons (Heisserer and Chambers 1993; Murray and Delahunty 2000a; Drake and others 2001b). Stampanoni (1994) identified food or chemical references for flavor lexicons for strawberry yogurts, caramel milk drinks, and cheese analogs to demonstrate standardization of sensory procedures and protocols. Civille and Lyon (1996) compiled a book focused on standardized definitions and references for a wide range of flavor descriptors. The use of references, particularly chemical references, since not all food types are available worldwide, allow for clarification of flavor lexicons for future use, and/or comparison. Fortunately, in recent years, peer review of sensory works has provided more intense pressure for definitions as well as references when publishing descriptive sensory research.

References can be qualitative, quantitative or both (Munoz and Civille 1998). For lexicons, qualitative references are critical for every term. Quantitative or intensity references are not generally provided for each attribute. Qualitative references allow panelists to associate with the concept of the term and can shorten panel training time (Rainey 1986). Qualitative references are used in training of panels to allow the panel to focus on the identified language and are a requisite part of most descriptive panel training. In contrast, fewer descriptive methods include quantitative references. Munoz and Civille (1998) described 3 types of quantitative references: universal, product specific, and attribute specific. These types of references also apply to 3 types of scaling: universal, product specific, and attribute specific.

Universal scale references cover the intensity range across all food categories. All attributes of any product are then rated rela-

tive to the universal scale intensity reference. An example would be the use of a 16% sucrose solution to demonstrate a 15 on the Spectrum universal scale. The solution demonstrates the intensity of the 15, and all tastes and flavors are rated accordingly. These references are external to the product category or the specific attribute. All attributes are rated using this same yardstick, so that attributes in a product are comparative across products, across attributes in 1 product, and over time and different panels. A product specific scale rates attributes within that specific product category. An intense flavor within that product category is assigned a high intensity. The upper end of the scale represents the highest intensities within the particular product category. In this manner, products within the category can be compared but cannot be compared to other products. In attribute specific scaling, each attribute is rated independently so that each attribute has its own intense quantitative reference. An example would be fruity and sulfur flavor in Cheddar cheese. Each attribute would have a high quantitative reference for that attribute within that product category. Fruity flavor is not an intense flavor in Cheddar cheese, while sulfur flavor is. The quantitative references for these 2 attributes on an attribute specific scale in Cheddar cheese would be quite different in intensity if compared to each other. With this scale, an attribute can be compared across products, but attributes cannot be compared within a product. Attribute specific scaling is closest to what untrained panelists and consumers do. It is also important to note that the 3 types of scaling probably represent the most to least amount of panelist training, with the universal scale requiring the most amount of training and attribute scaling, the least training.

A key benefit of a descriptive lexicon is its ability to relate to consumer acceptance/rejection and instrumental or physical measurements. A true "holy grail" lexicon would be one that was readily related to both consumer and instrumental measures. In reality, establishing even 1 relationship is a challenge, and often is not possible for some products. When key flavor terms for a product's category are represented by a group of chemicals that appear at low levels in the product (pyrazines and mercaptans for roasted coffee, peanuts, meat, chocolate), it is difficult to determine the exact mixture and amount of the various chemicals that characterize the "roasted" character. Also, no conclusions can be drawn regarding the nature of an attribute and whether the consumer response might be positive or not. For instance, the descriptor baby burp/vomit/butyric acid could all be used to accurately describe 1 particular flavor. Alternatively, baby burp/vomit could be used and butyric acid could be used as the reference. The danger would be in assuming any relation to consumer acceptance or rejection of this flavor in a particular product without extensive consumer testing. The use of consumer terminology in a descriptive language does not mean the language has direct application to consumer acceptability. The apparent "negative" connotation of vomit or butyric can be misleading, since butyric is an inherent and appropriate description for many cheeses and other fermented products. Further, consumer preferences and liking differ widely with age, geographic location, ethnic background, and familiarity with the product. What may constitute a "bad" flavor to one market segment may represent a "good" or "desirable" flavor to others.

In some cases, the lexicon has direct application to instrumental analysis. In the case of flavor lexicons, the lexicon has direct application to the volatile or nonvolatile compounds found in a product. Certainly, the use of chemical components, particularly chemical components isolated from that product, can make a lexicon more clear and grounded and can establish a link to formulation and/or production of that product. However, establishing those links can be a challenge and is addressed later in this review. Certainly, a lexicon can be representative, discriminative,

and precise without using consumer terminology or having chemical references.

Application of Specific Flavor Lexicons

Flavor lexicons are used widely to describe and discriminate among products within a category. They are widely used in industry for comparing and monitoring products and product consistency and for profiling new and competitive products. The use of lexicons in quality control, with relationships to instrumental or consumer responses, is full of fruitful applications. Lexicons are also a powerful research tool with numerous applications across many commodities and commercial products. Descriptive analysis has been conducted widely on wine. Noble and others (1984) first described a standardized and referenced language for wine aroma. The language has been widely applied, expanded, and built on and used throughout the wine industry for research (Heymann and Noble 1987; Noble and Shannon 1987; Noble and others 1987; McDaniel and others 1987; McCloskey and others 1996; Siversten and Risvik 1994).

In commodities such as cheese, several different flavor lexicons have been used to document aroma and flavor development, the effects of fat reduction, and the effects of different starter or adjunct bacteria. Muir and others (1995a) described 9 aroma terms for characterization of aroma profiles of hard and semi-hard cheese. Piggott and Mowat (1991) determined 23 descriptive flavor and aroma terms in a study of maturation of Cheddar cheese. McEwan and others (1989), Muir and Hunter (1992a), Muir and others (1996b) and Roberts and Vickers (1994) also developed descriptive flavor and aroma terms to study flavor during Cheddar cheese aging. Muir and others (1996a) and Drake and others (1996, 1997) used descriptive sensory panels to determine the effect of starter culture and adjunct cultures on Cheddar cheese flavor. Banks and others (1993) used descriptive analysis to determine sensory properties of low fat Cheddar cheese. A flavor lexicon for Comte cheese was used to identify naturally existing cheese georegions within France (Monnet and others 2000).

Sivertsen and Risvik (1994) demonstrated that French red wines could be differentiated according to region of origin using a sensory lexicon for wine. An aroma lexicon for meat/brothy aromas was used to optimize a meat-like process flavor made from enzyme hydrolyzed vegetable protein (Wu and others 2000). Several groups have identified and used descriptive sensory analysis to differentiate fluid milks (Lawless and Claassen 1993; Phillips and others 1995; Phillips and Barbano 1997; Watson and McEwan 1995; Chapman and others 2001; Bom Frost and others 2001). As might be expected, the languages have significant overlap in many terms. Flavor and aroma, as well as texture and feeling factors (aftertaste/mouthfeel) have been described. These lexicons have been used to determine the effects of fat content, storage, and other processing conditions on milk flavor and aroma perception. A flavor lexicon for chocolate ice cream was used to discern the effects of milkfat, cocoa butter, and fat replacers on sensory properties of chocolate ice creams (Prindiville and others 1999, 2000).

Descriptive analysis lexicons have also been used for numerous other products including almonds (Guerrero and others 1997), ripening anchovies (Bestiero and Rodriguez 2000), olive oil (Albi and Gutierrez 1991; Guerrero and others 2001) fermented milks (Hunter and Muir 1993; Muir and others 1997b), spoiled milk aroma (Hayes and others 2002), yogurt (Drake and others 2001a), beer (Meilgaard and others 1979), distilled beverages (Piggott and others 1980; Cristovam and others 2000; Lee and others 2001; McDonnell and others 2001) chicken (Lyon 1987), beef (Berry and others 1980), meat WOF (Johnsen and Cville 1986; Byrne and others 1999, 2001), Spanish cheeses (Ordonez and others

1998; Gonzalez-Vinas and others 1998; Barcenas and others 1999; Mendia and others 1999), cooked rice aroma (Yau and Liu 1999), milk chocolate (Mazzucchelli and Guinard 1999), catfish (Johnsen and others 1987), mutton (Rajalakshmi and others 1987), peanuts (Johnsen and others 1988), vanilla (Heymann 1994), ham (Flores and others 1997), apples (Dalliant-Spinnler and others 1996), processed cheese (Muir and others 1997a), rye bread (Hellemann and others 1987) sourdough bread (Lotong and others 2000). The languages can be very brief/concise or very descriptive. Cliff and Heymann (1991) used a language of 4 defined descriptors to describe the aroma of cat litter. Thirty seven descriptors were used to describe the aroma, flavor, and texture of dulce de leche (Hough and others 1992).

The value of lexicons in industry is underrepresented in technical publications because of the proprietary nature of the methods, lexicons, and resulting data. Most food retail companies use flavor lexicons throughout the development process. Early in the exploratory phase, R&D benchmarks the flavor characteristics of the potential competitive set and related products (Munoz and others 1995). During the development process, the descriptive panel uses the lexicon to document the development of different flavors in prototypes resulting from different ingredients, processes, and storage. At this phase, the lexicon often changes slightly to incorporate new attributes that develop in the new flavors. Once the final prototypes are ready for outside testing, the descriptive panel documents the products and prototypes going to the field test. Data relationships are established to determine the drivers of liking to set the best new product formulation and/or process. Once in production, these same attributes can be used to monitor production (Munoz and others 1992).

Benefit of References and Definitions in Lexicon Application

As previously mentioned, the use of definitions and references is important in establishing a language that can be reproduced at different sites or at different points in time. Previous research on descriptive sensory analysis of Cheddar cheese (Piggott and Mowat 1991; Roberts and Vickers 1994; Muir and Hunter 1992a; Muir and others 1995b) lacked definitions or references for descriptors. While some overlap between descriptors is observed, different research groups have used different descriptive vocabulary and rarely have definitions and references for descriptors been identified, making universal interpretation or reproduction of results difficult. A standard list of descriptors was identified for aged natural cheeses by Heisserer and Chambers (1993). More recently, Drake and others (2001b) identified a language for Cheddar cheese with standard definitions and references. Work is underway to confirm chemical references for this Cheddar cheese flavor lexicon. Murray and Delahunty (2000a) also developed and identified a referenced sensory lexicon for Cheddar-type cheeses. Both languages were developed independently, one in the U.S.A. (Drake and others 2001b) and one in Ireland (Murray and Delahunty 2000a). Studies have also been conducted to identify sensory attributes and descriptors for Parmesan and Comte cheeses (Virgili and others 1994; Berodier and others 1997).

Lexicons accurately communicate among research groups, particularly with a standardized lexicon and highly trained panelists. Previous studies with multiple panels at different sites have not used a uniform language. Risvik and others (1992) compared evaluation of chocolate between British and Norwegian panels, and Hirst and others (1994) later compared evaluation of cheese between British and Norwegian panels. Cross cultural differences were attributed to the observed discrepancies in term usage and sample differentiation. Sensory evaluation of hard cheese has

been continued in the European Union. Ring trials at 7 different sites across the EU were conducted, and a core sensory language for evaluation has been developed (Hunter and McEwan 1998; Nielson and Zannoni 1998). However, definitions and references (anchors) for each descriptor were not identified. Lundgren and others (1986) conducted an interlaboratory international study on firmness, aroma intensity, sweetness, sourness, and flavor intensity of pectin gels using untrained panelists, and noted good agreement of the results, indicating that simple terms in simple systems may be universal in perception. Lotong and others (2001) reported the results of descriptive analysis of orange juice flavor by 2 highly trained descriptive panels using different descriptive analysis methods and independent lexicons. Similar patterns of differentiation among samples were observed between the 2 groups. Pages and Husson (2001) also studied panel performance with 13 different groups for 2 foods: compotes and chocolate. Efforts were made to standardize the descriptors and scale among the groups prior to the study, although the amount of panel training for each group was not specified. Again, similar patterns of differentiation were observed between the panels, although there were minor differences in scale usage and descriptor significance.

While similar patterns of differentiation among samples by panels that use different languages are expected (particularly if the lexicons are comprehensive and the panelists highly trained), standardized language with definitions and references improve communication, cross panel validation, and subsequent application of descriptive analysis results to instrumental or consumer data. Further, other sources of variation potentially exist in comparing panel results at different sites within the same country using the same language. Drake and others (2002) reported on the performance of 3 descriptive panels trained at different sites by different panel leaders on a previously developed and standardized cheese lexicon (Drake and others 2001b). Panels were able to accurately communicate attribute differences between cheeses. There were observed differences between these panels in scale usage and attribute recognition. These differences were attributed to differences in panel leadership and h of panelist training. In a similar study, Martin and others (2000) compared odor profile results of 2 panels. Language, scale, and method of presentation were standardized. Results obtained from the 2 panels were similar. Differences between attribute intensities were reported and were attributed to differences in individual panelist experience and/or perception. As with the conclusions of Drake and others (2002), strong panel leader interaction was recommended as a method to rectify these differences, along with regular feedback between the 2 panels.

Relating Sensory Perception to Consumers

While descriptive sensory analysis is used to identify and quantify information on the sensory aspects of products, effective consumer tests are used to provide information on consumer liking (Meilgaard and others 1999). Acceptability and preference tests provide quantitative consumer liking and/or preference information. Screeners and questionnaires are used to gather demographic data, frequency of usage, and purchase history about a particular product or group of products (Lawless and Heymann 1998; Meilgaard and others 1999). Questionnaires are often included with and are recommended with acceptance testing to aid in data interpretation. Despite collection of demographic and usage information, identifying drivers of liking or disliking within consumer market segments is critical for industry to identify which products and product attributes are preferred and by which consumer market segment.

In order to relate consumer liking and descriptive sensory properties, several statistical methods are applied. Simple techniques,

such as scatter plots and correlation analysis, are critical to learning about the possible direct relationships between sensory attributes and consumer liking. Often, multivariate analysis is required to fully probe and relate the data sets and determine the combinations of attributes that affect liking. Multivariate techniques used include multiple linear regression analysis, canonical correlation analysis, principal component analysis, partial least squares regression analysis, and preference mapping, a principal component analysis derivation (Munoz and Chambers 1993; Heyd and Danzart 1998; Barcenas and others 2001). A group of products are documented with descriptive analysis and then tested with consumers. The analytical descriptive properties, which describe exactly what attributes are perceived and at what levels, are related to the consumer preferences using combinations of multivariate techniques. "Preference mapping", as this relationship building is called, has been conducted with many products including cheese (McEwan and others 1989, Lawlor and Delahunty 2000; Murray and Delahunty 2000b; Barcenas and others 2001), chicken nuggets (Arditti 1997), coffee (Heyd and Danzart 1998), apples (Dalliant-Spinnler and others 1996; Jaeger and others 1998), fermented milks (Muir and Hunter 1992b), sheepmeat (Prescott and others 2001), pudding (Elmore and others 1999), milks of varying fat content (Richardson-Harman and others 2000), and cured meats (Munoz and others 1995).

One of the biggest challenges in consumer research is the clarification of consumer language. Consumers may use terms that are ambiguous, have multiple meanings, are associated with "good" or "bad", or are combinations of several terms. These integrated terms, such as "creamy", are often used by consumers to represent a combination of positive attributes. Determining exactly what attribute or attributes "creamy" refers to (flavor compared with texture compared with mouthfeel) has been the subject of many studies relating consumer and trained sensory panels to clarify this term (Mela 1988; Elmore and others 1999; Bom Frost and others 2001). Dacremont and Vickers (1994a) used concept matching to clarify consumer perception of Cheddar cheese flavor. Descriptive analysis was not conducted to correlate with the obtained consumer information, and a limited number of consumers ($n = 18$) were used for the assessments. The concept of Cheddar cheese flavor is a consumer concept and likely varies widely among consumers, as does Cheddar cheese flavor itself. A subsequent study was conducted by the authors using combinations of chemical volatiles (isolated from cheese) and concept matching to further clarify the role of individual and combinations of chemicals on the Cheddar cheese flavor concept (Dacremont and Vickers 1994b). Again, a limited number of subjects were used in the flavor concept experiments ($n = 16$ to 21), which potentially limits conclusions from the study. Doubtless, with large experiments and multiple combinations of treatments, the use of large numbers of consumers is impractical, but care must be taken when interpreting such data. Further studies with larger consumer groups and demographic information including types (brand, age) of Cheddar cheese normally consumed would provide additional clarification.

Relating Sensory Perception to Chemical Components

Relating sensory language and chemical volatile compounds represents a challenge for several reasons. Relative amount of a compound in a food is not necessarily a measure of its sensory impact, due to different thresholds and the effects of the food matrix. The sensitivity of the extraction technique must also be taken into account. Finally, only a small percentage of the volatile components in a food are odor-active (Friedrich and Acree 1998). The 1st step in the solution to relate gas chromatographic data and chemical compounds to sensory impact is gas-chromatography

olfactometry (GC/O). In this technique, the individual compounds in the GC effluent are sniffed and described by a trained panelist. The technique can also be quantitative if the relative intensity of the odorant may also be recorded. Three different techniques are commonly applied with GC/O: aroma extract dilution analysis (AEDA), CharmAnalysis™, and OSME. These techniques and others have been loosely grouped into 3 categories: dilution analysis, posterior intensity methods, and detection frequency methods (van Ruth and O'Connor 2001). Both AEDA and CharmAnalysis operate on dilution of samples until an odor is no longer detected. The highest dilution at which the odor is detected can be converted to a flavor dilution value (FD value used with AEDA) or a Charm value (Friedrich and Acree 1998). The requisite sniffing of large numbers of sample dilutions by at least 2 panelists makes these techniques time-consuming. These techniques also assume that the response to a stimulus is linear and that all compounds have the same linear response. Osme, in contrast, is a time intensity technique and does not involve dilutions, but instead requires panelists (3 or more) to evaluate the time-intensity of aromas as well as the aroma character (Miranda-Lopez and others 1992; Plotto and others 1998; Fu and others 2002). Posterior intensity techniques can also involve 2 or more trained panelists simply scoring the maximum perceived intensity of an aroma along with its aroma character. A recent olfactometry technique, nasal impact frequency (NIF) has also been developed based on frequency of detection of a compound in the GC effluent by sensory panelists (Pollien and others 1997; Bezman and others 2001). The NIF technique can be applied with time intensity to yield surface of nasal impact frequency (SNIF) data. Pollien and others (1999) demonstrated that NIF and SNIF could be applied to quantify 1-octen-3-one in solution and in coffee aroma with results comparable to mass spectrometry. Both Osme and other posterior intensity methods and NIF/SNIF require fewer GC runs than AEDA and CharmAnalysis, since only the undiluted flavor extract is evaluated by the sensory panelist. However, a larger number of panelists (3 or more, compared to 2 for AEDA and CharmAnalysis) is recommended for these techniques. All of these techniques are useful for determining odor activity of volatile compounds.

One of the limitations to these techniques is using few panelists due to the time requirement. Usually a minimum of 2 sniffers is advised. Finally, the information obtained is based on individual components and does not include matrix effects or the time release factor involved when a sample is chewed in the mouth. Another device, the artificial mouth or RAS (retronasal aroma simulation) addresses the conditions encountered by food in the mouth (Roberts and Acree 1996; Friedrich and Acree 1998). Saliva, pH, temperature, and mixing can all be addressed, followed by subsequent analysis of the released volatiles in the headspace. Van Ruth and others (2001) studied the effects of saliva components of volatile release of 20 aroma compounds in model systems by static headspace analysis. Their goal was to relate results to volatile flavor release in the mouth. Their findings suggested that, while saliva composition does influence volatile release, the ratio of saliva to matrix played a larger role. Flavor release and perception is a very complex process (Taylor 1996). RAS simplifies a very complex process, and, while valuable insight is gained with this technique, the time release aspect of food consumption and subsequent flavor detection and the matrix effects are still not addressed.

Numerous studies beyond the scope of this review have been conducted on aroma volatiles of selected foods with GC and GC/O techniques. For the reasons mentioned previously, these results are useful but limited in any application to sensory perception (Piggott 1990). A list of volatile components with their odor properties does not equal sensory flavor. Descriptive analysis of the food product(s) can be done simultaneously with the instrumental

analysis and the results analyzed for statistical relationships by univariate and multivariate analysis of variance. Karagul-Yuceer and others (2001, 2002) conducted instrumental analysis (including GC/O) and descriptive sensory analysis of nonfat dried milk powders to determine potential relationships between instrumental and sensory perceptions of flavor. Cadwallader and Howard (1998) conducted a similar study of light activated milk flavor using instrumental analyses and descriptive sensory analysis, as did Young and others (1996) with apple flavor and Maeztu and others (2001) with espresso coffee aroma. When a control product and a product identical except for the flavor of interest is available, a side-by-side comparison can also be conducted. Suriyaphan and others (1999) identified alkadienals responsible for off-flavors in cheeses containing lecithin by a side-by-side instrumental comparison of experimental cheeses made with and without added lecithin. While potential relationships can be identified in this manner, subsequent work is required to further establish these links. The subsequent steps in relating instrumental data to sensory perception in a food involve sensory analysis, threshold analysis, and the use of model systems or the addition of a suspected flavor compound directly to the food.

Sensory threshold data can expand understanding of odor active data from GC/O. The ratio of the concentration of an odorant to its odor threshold is expressed as odor active value (OAV). OAV's are often considered a more accurate determination of a compound's contribution to perceived flavor (Preininger and Grosch 1994; Friedrich and Acree 1998). Thus, sensory threshold analysis is often conducted on key aroma active compounds identified and quantified by instrumental analysis. The OAV's are then determined and evaluated for significance prior to model system studies. Addition of the compounds to a bland matrix of the particular food or a model system followed by descriptive analysis provides further links to the actual role of the compound in human perception of flavor. Belford and others (1991) used descriptive analysis with a model syrup system with selected added volatiles to determine important volatile components of maple syrup. Guth and Grosch (1994) identified and quantified potent odorants from stewed beef juice by aroma extract dilution analysis. Odorants were then incorporated into a model (odor-free) beef juice system, into which selected key odorants were added. The beef juice models were then evaluated by a trained panel for their similarity to stewed beef juice using difference-from-control testing, with stewed beef juice as the control. Odorants were removed from the model mixtures sequentially and the similarity to stewed beef juice assessed. In this manner, contributing odorants were identified. Similar studies using model systems have been applied for strawberry juice, Gala apple flavor, meat-like process flavoring, virgin olive oil, orange juice, and sourdough rye bread (Schieberle and Hofman 1997; Plotto and others 1998; Wu and others 2000; Reiners and Grosch 1998; Buettner and Schieberle 2001a, 2001b; Kirchoff and Schieberle 2001). Compounds can also contribute to flavor at concentrations at or below sensory threshold, due to interactions with other chemicals or the food matrix. Unfortunately, the role of these compounds is not addressed with this technique.

Panelists tasted water-soluble extracts of comte cheese to identify fractions which had particular tastes in an attempt to clarify the role of peptides and amino acids on flavor (Salles and others 1995). Preininger and others (1996) used an unripened cheese matrix to evaluate both volatiles and nonvolatile flavor components of 2 Swiss cheese samples. A similar study was conducted with Emmental cheese (Rychlik and others 1997). Suriyaphan and others (2001) identified key chemical volatile components of cowy/barny and earth/bell pepper sensory perceived flavors in selected aged British Farmhouse cheeses. In this study, sensory properties were identified by descriptive sensory analysis, aroma

volatiles were quantified by GC/MS, and aroma properties described by GC/O. Suspected key volatiles were selected from GC/O data based on aroma properties and FD values. The selected aroma components were subsequently incorporated into mild (bland) cheese across the concentration range encountered in the Farmhouse cheeses and evaluated by descriptive analysis. Studies such as these provide convincing evidence of the role of particular compounds on flavor. Model systems have not as yet provided insights into the role of compound mixtures and the role of compounds at sub-threshold.

Conclusion

Flavor lexicons are an indispensable tool for accurately documenting the description of food flavor. As with any scientific method, sensory scientists need to adhere to those recommended practices required to assure panel performance: careful screening and selection of panelists; comprehensive training that includes lexicon development; definitive references; scaling standardization; experimental controls for sample preparation and evaluation; and ongoing maintenance of the panel system. Future research will continue to address relating descriptive (and analytical) lexicons with consumer acceptance and instrumental flavor analysis. More testing is expected and needed to demonstrate the robustness of the methodology and its practical application across cultures, languages, descriptive methods, and analytical chemical techniques.

References

- Albi MA, Gutierrez F. 1991. Study of the precision of an analytical taste panel for sensory evaluation of virgin olive oil. Establishment of criteria for the elimination of abnormal results. *J Sci Food Agric* 54:225-67.
- Arditti S. 1997. Preference mapping: a case study. *Food Qual. Pref.* 8:323-7.
- Banks JM, Hunter EA, Muir DD. 1993. Sensory properties of low fat Cheddar: effect of salt content and adjunct culture. *J Soc Dairy Technol* 46:119-23.
- Barcenas P, Elortondo FJP, Salmeron J, Albisu M. 1999. Development of a preliminary sensory lexicon and standard references of ewes milk cheeses aided by multivariate statistical procedures. *J Sensory Stud* 14:161-79.
- Barcenas P, San Roman RP, Elortondo FJP, Albisu M. 2001. Consumer preference structures for traditional Spanish cheeses and their relationship with sensory properties. *Food Qual Pref* 12:269-79.
- Belford AL, Lindsay RC, Ridley SC. 1991. Contributions of selected flavor compounds to the sensory properties of maple syrup. *J Sens Stud* 6:101-18.
- Berodier F, Stevenot C, Schlich P. 1997. Descripteurs de l'arome du fromage de Comte. *Lebensm Wiss Technol* 30:298-304.
- Berry BW, Maga JA, Calkins CR, Wells LH, Carpenter AL, Cross HR. 1980. Flavor profile analysis of cooked beef loin steaks. *J Food Sci* 45:1113-5.
- Bestiero I, Rodriguez CJ. 2000. Selection of attributes for the sensory evaluation of anchovies during the ripening process. *J Sens Stud* 15:65-77.
- Bezman Y, Rousset RL, Naim M. 2001. 2-methyl-3-furanthiol and methional are possible off-flavors in stored orange juice: aroma-similarity, NIF/SNIF GC-O, and GC analyses. *J Agric Food Chem* 49:5425-32.
- Bom Frost M, Dijksterhuis GB, Martens, M. 2001. Sensory perception of fat in milk. *Food Qual Pref* 12:327-36.
- Buettner A, Schieberle P. 2001a. Evaluation of aroma differences between hand-squeezed juices from Valencia late and Navel oranges by quantitation of key odorants and flavor reconstitution experiments. *J Agric Food Chem* 49:2387-94.
- Buettner A, Schieberle P. 2001b. Evaluation of Key Aroma Compounds in Hand-Squeezed Grapefruit Juice (*Citrus paradisi* Macfayden) by Quantitation and Flavor Reconstitution Experiments. *J Agric Food Chem* 49:1358-63.
- Byrne DV, Bak LS, Bredie WLP, Bertelson G, Martens M. 1999. Development of a sensory vocabulary for warmed-over flavor: Part 1. In porcine meat. *J Sensory Stud* 14:47-65.
- Byrne DV, O'Sullivan MG, Dijksterhuis GB, Bredie WLP, Martens M. 2001. Sensory panel consistency during development of a vocabulary for warmed-over flavor. *Food Qual Pref* 12:171-87.
- Cadwallader KR, Howard CL. 1998. Analysis of aroma active components of light activated milk. In: Mussinan CJ, Morello MJ, editors. *Flavor Analysis, Developments in Isolation and Characterization*. ACS symposium Series. Washington DC: American Chemical Society. p 343-58.
- Chapman KW, Lawless HT, Boor KJ. 2001. Quantitative descriptive analysis and principal component analysis for sensory characterization of ultrapasteurized milk. *J Dairy Sci* 84:12-20.
- Civille GV, Lawless HT. 1986. The importance of language in describing perceptions. *J Sens Stud* 1:203-15.
- Civille GV, Lyon BG. 1996. Aroma and Flavor Lexicons for Sensory Evaluation: Terms, Definitions, and References. ASTM D5 66, West Conshohocken, Pa: Am Soc Testing and Materials. 158 p.
- Cliff M, Heymann, H. 1991. Physical and sensory characteristics of cat litter. *J Sens Stud* 6:255-66.

- Cristovam E, Paterson A., Piggott JR. 2000. Development of a vocabulary of terms for sensory evaluation of dessert port wines. *Ital J Food Sci* 12:129-42.
- Dacremont C, Vickers Z. 1994a. Classification of cheeses according to their closeness to the Cheddar cheese concept. *J Sens Stud* 9:237-246.
- Dacremont C, Vickers Z. 1994b. Concept matching techniques for assessing importance of volatile compounds for Cheddar cheese aroma. *J Food Sci* 59:981-5.
- Dailliant-Spinner B, Macfie HJH, Beyts PK, Hedderley D. 1996. Relationships between perceived sensory properties and major preference directions of 12 varieties of apples from the southern hemisphere. *Food Qual Pref* 7:113-26.
- Drake MA, Boylston TD, Spence KD, Swanson BG. 1996. Chemical and Sensory characteristics of reduced fat cheese with a *Lactobacillus* adjunct. *Food Res Int* 29(3/4):381-7.
- Drake MA, Boylston TD, Spence KD, Swanson BG. 1997. Improvement of sensory quality of reduced fat Cheddar cheese by a *Lactobacillus* adjunct. *Food Res Int* 30(1):35-40.
- Drake MA, Gerard PD, Chen XQ. 2001a. Effects of sweetener, sweetener concentration, and fruit flavor on sensory properties of soy fortified yogurt. *J Sens Stud* 16:393-406.
- Drake MA, McIngvale SC, Gerard PD, Cadwallader KR, Civiile GV. 2001b. Development of a descriptive language for Cheddar cheese. *J Food Sci* 66:1422-1427.
- Drake MA, Gerard PD, Wright S, Cadwallader KR, Civiile GV. 2002. Cross validation of a sensory language for Cheddar cheese. *J Sens Stud* 17:215-29.
- Elmore JR, Heymann H, Johnson J, Hewett JE. 1999. Preference odorings: relating acceptance of "creaminess" to a descriptive sensory map of a semi-solid. *Food Qual Pref* 10:465-75.
- Flores M, Ingram DA, Bett KL, Toldra F, Spanier AM. 1997. Sensory characteristics of Spanish Serrano dry cured ham. *J Sens Stud* 12:169-79.
- Friedrich JE, Acree TE. 1998. Gas chromatography olfactometry (GC/O) of dairy products. *Int Dairy J* 8:235-41.
- Fu S, Yoon Y, Bazemore R. 2002. Aroma-active components of fermented bamboo shoots. *J Agric Food Chem* 50:549-54.
- Gonzalez Vinas MA, Esteban EM, Cabezas L. 1998. Physico-chemical and sensory properties of Spanish ewe milk cheeses and consumer preferences. *Milchwissenschaft* 54:326-9.
- Guerrero L, Gou P, Arnau J. 1997. Descriptive analysis of toasted almonds: a comparison between experts and semi-trained assessors. *J Sens Stud* 12:39-54.
- Guerrero L, Romero A, Tous J. 2001. Importance of generalized procrustes analysis in sensory characterization of virgin olive oil. *Food Qual Pref* 12:515-20.
- Guth H, Grosch W. 1994. Identification of the characteristic impact odorants of stewed beef juice by instrumental analysis and sensory studies. *J Agric Food Chem* 42:2862-6.
- Hayes WW, White CH, Drake MA. 2002. Sensory aroma characteristics of milk spoilage by *Pseudomonas* species. *J Food Sci* 67:861-7.
- Heisserer DM, Chambers E. 1993. Determination of the sensory flavor attributes of aged natural cheese. *J Sens Stud* 8:121-32.
- Hellermann U, Tuorila H, Salovaara H, Tarkkonen L. 1987. Sensory profiling and multidimensional scaling of Finnish rye breads. *Int J Food Sci Technol* 22:693-700.
- Heyd B, Danzart M. 1998. Modelling consumers' preferences of coffees: evaluation of different methods. *Lebensm Wiss Technol* 31:607-611.
- Heymann H, Noble AC. 1987. Descriptive analysis of commercial cabernet sauvignon wines from California. *Am J Enol Vitic* 38:41-4.
- Heymann H. 1994. A comparison of descriptive analysis of vanilla by 2 independently trained panels. *J Sens Stud* 9:21-32.
- Hirst D, Muir DD, Naes, T. 1994. Definition of the sensory properties of hard cheese: a collaborative study between Scottish and Norwegian Panels. *Int Dairy J* 4:743-61.
- Hough G, Bratchell N, Macdougall DB. 1992. Sensory profiling of dulce de leche, a dairy based confectionary product. *J Sens Stud* 7:157-78.
- Hunter EA, Muir DD. 1993. Sensory properties of fermented milks: objective reduction of an extensive sensory vocabulary. *J Sens Stud* 8:213-27.
- Hunter EA, McEwan JA. 1998. Evaluation of an international ring trial for sensory profiling of hard cheese. *Food Qual Pref* 9:343-54.
- Ishii R, O'Mahoney M. 1987. Defining a taste by a single standard: aspects of salty and umami tastes. *J Food Sci* 52:1405-9.
- Ishii R, O'Mahoney M. 1991. Use of multiple standards to define sensory characteristics for descriptive analysis: aspects of concept formation. *J Food Sci* 56:838-42.
- Jaeger SR, Andani Z, Wakeling IN, Macfie HJH. 1998. Consumer preferences for fresh and aged apples: a cross cultural comparison. *Food Qual Pref* 9:355-66.
- Janto M, Pipatsattayanuwong S, Kruk MW, Hou G, McDaniel MR. 1998. Developing noodles from US wheat varieties for the far east market: sensory perspective. *Food Qual Pref* 9:403-12.
- Johnsen PB. 1986. A lexicon of peanut flavor descriptors. ARS-52 Washington DC: US Dept. of Agriculture. 8 p.
- Johnsen PB, Civiile GV. 1986. A standardized lexicon of meat WOF descriptors. *J Sens Stud* 1:99-104.
- Johnsen PB, Civiile GV, Vercellotti JR. 1987. A lexicon of pond-raised catfish flavor descriptors. *J Sens Stud* 2:85-91.
- Johnsen PB, Civiile GV, Vercellotti J, Sanders TH, Dus CA. 1988. Development of a lexicon for the description of peanut flavor. *J Sens Stud* 3:9-18.
- Karagul-Yuceer Y, Drake MA, Cadwallader KR. 2001. Aroma-active components of nonfat dried milk. *J Agric Food Chem* 49:2948-53.
- Karagul-Yuceer Y, Drake MA, Cadwallader KR. 2002. Volatile flavor components of stored nonfat dry milk. *J Agric Food Chem* 50:305-12.
- Keane P. 1992. The flavor profile. In: R.C. Hootman, editor. ASTM Manual Series MNL 13 Manual on Descriptive Analysis Testing for Sensory Evaluation. West Conshohocken, PA: Am. Soc. Testing and Materials. p 5-14.
- Kirchhoff E, Schieberle P. 2001. Determination of key aroma compounds in the crumb of a 3-stage sourdough rye bread by stable isotope dilution assays and sensory studies. *J Agric Food Chem* 49:4304-11.
- Lawless HT, Claassen MR. 1993. Validity of descriptive and defect-oriented terminology systems for sensory analysis of fluid milk. *J Food Sci* 58:108-12,119.
- Lawless HT, Heymann H. 1998. Qualitative consumer research methods. In: Sensory Evaluation of Food. New York NY: Chapman and Hall. p 519-47.
- Lawlor JB, Delahunty CM. 2000. The sensory profile and consumer preference for ten specialty cheeses. *Int J Dairy Technol* 53:28-36.
- Lee KY M, Paterson A, Piggott JR, Richardson, GD. 2001. Sensory discrimination of blended Scotch whiskies of different product categories. *Food Qual Pref* 12:109-17.
- Lotong V, Chambers E, Chambers DH. 2000. Determination of the sensory attributes of wheat sourdough bread. *J Sens Stud* 15:309-26.
- Lotong V, Chambers DH, Dus C, Chambers E, Civiile, GV. 2001. Matching results of 2 independent highly trained panels using different descriptive analysis methods [abstract]. In: 4th Pangbor Sensory Science Symposium; Dijon, France; July 22-26. abstract P-060.
- Lundgren B, Pangborn RM, Daget N, Yoshida M, Laing DG, McBride RL, Griffeths N, Hyvonen L, Sauvageot F, Paulus K, Barylko-Pikielna N. 1986. An interlaboratory study of firmness, aroma, and taste of pectin gels. *Lebensm Wiss Technol* 19:66-76.
- Lyon BG. 1987. Development of chicken flavor descriptive attribute terms aided by multivariate statistical procedures. *J Sens Stud* 2:55-67.
- Maezta L, Sanz C, Andueza S, Paz de Pena M, Bello J, Cid C. 2001. Characterization of espresso coffee aroma by static headspace GC-MS and sensory flavor profile. *J Agric Food Chem* 49:5437-44.
- Martin N, Molimard P, Spinner HE, Schlich P. 2000. Comparison of odour sensory profiles performed by 2 independent trained panels following the same descriptive analysis procedures. *Food Qual Pref* 11:487-95.
- Mazzucchelli R, Guinard JX. 1999. Comparison of monadic and simultaneous sample presentation modes in a descriptive analysis of milk chocolate. *J Sens Stud* 14:235-48.
- McCloskey LP, Sylvan M, Arrhenius SP. 1996. Descriptive analysis for wine quality experts determining appellations by chardonnay wine aroma. *J Sens Stud* 11:49-67.
- McDaniel M, Henderson LA, Watson BT, Heatherbell D. 1987. Sensory panel training and screening for descriptive analysis of the aroma of pinot noir wine fermented by several strains of malolactic bacteria. *J Sens Stud* 2:149-67.
- McDonnell E, Hulin-Bertaud S, Sheehan EM, Delahunty CM. 2001. Development and learning process of a sensory vocabulary for the odor evaluation of selected distilled beverages using descriptive analysis. *J Sens Stud* 16:425-45.
- McEwan JA, Moore JD, Colwill JS. 1989. The sensory properties of Cheddar cheese and their relationship with acceptability. *J Soc Dairy Technol* 42:112-7.
- Manual on Descriptive Analysis Testing for Sensory Evaluation [M.D.A.T.S.E.]. 1992. ASTM manual series MNL 13. Hootman RC, editor. West Conshohocken, Pa.: American Society for Testing and Materials. 52 p.
- Meilgaard MM, Civiile GV, Carr, T. 1999. Sensory Evaluation Techniques. 3rd Ed. New York, NY: CRC Press. 387 p.
- Meilgaard MC, Daigliesh CE, Clapperton JF. 1979. Beer flavour terminology. *J Inst Brew* 85:38-42.
- Meilgaard MC, Reid DS, Wyborski KA. 1982. Reference standards for beer flavor terminology system. *Am Soc Brew Chem J* 40:119-28.
- Mela DJ. 1988. Sensory assessment of fat content in fluid dairy products. *Appetite* 10:37-44.
- Mendia C, Ibanez FC, Torre P, Barcina Y. 1999. Effect of pasteurization on the sensory characteristics of ewe's milk cheese. *J Sens Stud* 14:415-24.
- Miranda-Lopez R, Libbey LM, Watson B, McDaniel MR. 1992. Odor analysis of Pinot Noir wines from grapes of different maturities by a gas chromatography-olfactometry technique (Osme). *J Food Sci* 57:985-93, 1019.
- Monnet JC, Berodier F, Badot PM. 2000. Characterization and localization of a cheese region using edaphic criteria (Jura Mountains, France). *J Dairy Sci* 83:1692-1704.
- Muir DD, Hunter EA. 1992a. Sensory evaluation of Cheddar cheese: the relation of sensory properties to perception of maturity. *J Soc Dairy Technol* 45:23-29.
- Muir DD, Hunter EA. 1992b. Sensory evaluation of fermented milks: vocabulary development and the relations between sensory properties and composition and between acceptability and sensory properties. *J Soc Dairy Technol* 45:73-80.
- Muir DD, Hunter EA, Watson M. 1995a. Aroma of cheese. 1. Sensory characterization. *Milch* 50:499-503.
- Muir DD, Hunter EA, Banks JM, Horne DS. 1995b. Sensory properties of hard cheese: identification of key attributes. *Int Dairy J* 5:157-77.
- Muir DD, Banks JM, Hunter EA. 1996a. Sensory properties of Cheddar cheese: effect of starter type and adjunct. *Int Dairy J* 6:407-23.
- Muir DD, Hunter EA, Banks JM, Horne DS. 1996b. Sensory properties of Cheddar cheese: changes during maturation. *Food Res Int* 28(6):561-8.
- Muir DD, Williams SAR, Tamime AY, Shenana ME. 1997a. Comparison of the sensory profiles of regular and reduced fat commercial processed cheese spreads. *Int J Food Sci Technol* 32:279-87.
- Muir DD, Hunter EA, Dalaudier C. 1997b. Association of the sensory properties of commercial, strawberry flavoured fermented milks with product composition. *Int J Dairy Technol* 50:28-34.
- Munoz AM, Civiile GV. 1992. Spectrum descriptive analysis. In: ASTM Manual Series MNL 13 Manual on Descriptive Analysis Testing for Sensory Evaluation. Hootman RC, editor, West Conshohocken, Pa.: Am. Soc. Testing and Materials. p 22-34.
- Munoz AM, Chambers E, Hummer SA. 1995. Multifaceted category research study: how to understand a product category and its consumer responses. *J Sens Stud* 11:261-94.
- Munoz AM, Civiile GV. 1998. Universal, product, and attribute specific scaling and the development of common lexicons in descriptive analysis. *J Sens Stud* 13:57-76.
- Munoz AM, Civiile GV, Carr, BT. 1992. Sensory Evaluation in Quality Control. Chatam, N.J.: Sensory Spectrum, Inc. 240 p.
- Munoz AM, Chambers E. 1993. Relating sensory measurements to consumer acceptance of meat products. *Food Technol* 47:128-31, 134.
- Murray JM, Delahunty CM. 2000a. Selection of standards to reference terms in a Cheddar-type cheese flavor language. *J Sens Stud* 15:179-99.
- Murray JM, Delahunty CM. 2000b. Mapping consumer preferences for the sensory and packaging attributes of Cheddar cheese. *Food Qual Pref* 11:419-35.
- Nielson RG, Zannoni M. 1998. Progress in developing an international protocol for sensory profiling of hard cheese. *Int J Dairy Technol* 51:57-64.
- Noble AC, Arnold RA, Masuda BM, Pecore SD, Schmidt JO, Stern PM. 1984. Progress towards a standardized system of wine aroma terminology. *Am J Enol Vitic* 35:107-9.
- Noble AC, Shannon M. 1987. Profiling zinfandel wines by sensory and chemical

- analyses. *Am J Enol Vitic* 38:1-5.
- Noble AC, Arnold J, Buechsenstein A, Leach EJ, Schmidt JO, Stern PM. 1987. Modification of a standardized system of wine aroma terminology. *Am J Enol Vitic* 38:143-6.
- Ordóñez AI, Ibanez FC, Torre P, Barcina Y. 1998. Application of multivariate analysis to sensory characterization of ewe's milk cheese. *J Sens Stud* 13:45-55.
- Pages J, Husson, F. 2001. Inter-laboratory comparison of sensory profiles: methodology and results. *Food Qual Pref* 12:297-309.
- Pangborn RM. 1989. The evolution of sensory science and its interaction with IFT. *Food Technol* 43:248-56, 307.
- Phillips LG, McGiff M, Barbano DM, Lawless HT. 1995. The influence of fat on the sensory properties, viscosity, and color of lowfat milk. *J Dairy Sci* 78:1258-66.
- Phillips LG, Barbano DM. 1997. The influence of fat substitutes based on protein and titanium dioxide on the sensory properties of lowfat milks. *J Dairy Sci* 80:2726-31.
- Piggott JR, Hose LP, Sharp R. 1980. Expressing flavor: a system for scotch. *Brew Distil Int* 10:48-50.
- Piggott JR. 1990. Relating sensory and chemical data to understand flavor. *J Sens Stud* 4:261-72.
- Piggott JR, Mowat RG. 1991. Sensory aspects of maturation of Cheddar cheese by descriptive analysis. *J Sens Stud* 6:49-62.
- Piggott JR, Simpson SJ, Williams SAR. 1998. Sensory analysis. *Int J Food Sci Technol* 33:7-18.
- Plotto A, Mattheis JP, Lundahl DS, McDaniel MR. 1998. Validation of gas chromatography olfactometry results for Gala apples by evaluation of aroma-active compound mixtures. In: Mussinan CJ, Morello MJ, editors. *Flavor Analysis, Developments in Isolation and Characterization*. Washington DC: ACS symposium Series, American Chemical Society, p 290-302.
- Pollien P, Ott A, Montigton F, Baumgartner M, Munoz-Box R, Chaintreau A. 1997. Hyphenated headspace gas chromatography sniffing technique: screening of impact odorants and quantitative aromagram comparisons. *J Agric Food Chem* 4:2630-7.
- Pollien P, Fay LB, Baumgartner M, Chaintreau A. 1999. First attempt at odorant quantitation using gas chromatography-olfactometry. *Anal Chem* 71:5391-7.
- Preininger M, Grosch W. 1994. Evaluation of key odorants of the neutral volatiles of Emmentaler cheese by the calculation of odour activity values. *Lebensm Wiss Technol* 27:237-43.
- Preininger M, Warmke R, Grosch W. 1996. Identification of the character impact flavour compounds of Swiss cheese by sensory studies of models. *Z Lebensm Unters Forsch* 202:30-4.
- Prescott J, Young O, O'Neill L. 2001. The impact of variations in flavour compounds on meat acceptability: a comparison of Japanese and New Zealand consumers. *Food Qual Pref* 12:257-64.
- Prindiville EA, Marshall RT, Heymann H. 1999. Effect of milk fat on the sensory properties of chocolate ice cream. *J Dairy Sci* 82:1425-32.
- Prindiville EA, Marshall RT, Heymann H. 2000. Effect of milk fat, cocoa butter, and whey protein fat replacers on the sensory properties of lowfat and nonfat chocolate ice cream. *J Dairy Sci* 83:2216-23.
- Rainey B. 1986. Importance of reference standards in training panelists. *J Sens Stud* 1:149-54.
- Rajalakshmi D, Dhanaraj S, Chand N, Govindarajan VS. 1987. Descriptive quality analysis of mutton. *J Sens Stud* 2:93-118.
- Reiners J, Grosch W. 1998. Odorants of virgin olive oils with different flavor profiles. *J Agric Food Chem* 46:2754-63.
- Richardson-Harman NJ, Stevens R, Walker S, Gamble J, Miller M, Wong M, McPherson A. 2000. Mapping consumer perceptions of creaminess and liking for liquid dairy products. *Food Qual Pref* 11:239-46.
- Risvik E, Colwill JS, McEwan JA, Lyon DH. 1992. Multivariate analysis of conventional profiling data: a comparison of a British and a Norwegian panel. *J Sens Stud* 7:97-118.
- Roberts AK, Vickers ZM. 1994. Cheddar cheese aging: changes in sensory attributes and consumer acceptance. *J Food Sci* 59:328-34.
- Roberts DD, Acree TE. 1996. Effects of heating and cream addition on fresh raspberry aroma using a retronasal aroma simulator and gas chromatography olfactometry. *J Agric Food Chem* 44:3919-25.
- Rychlik M, Warmke R, Grosch W. 1997. Ripening of Emmentaler cheese wrapped in foil with and without addition of *Lactobacillus casei* ssp. *casei*. III. Analysis of character impact flavour compounds. *Lebensm Wiss Technol* 30:471-8.
- Salles C, Septier C, Roudot-Algaron F, Guillot A, Etievant PX. 1995. Sensory and chemical analysis of fractions obtained by gel permeation of water soluble comte cheese extracts. *J Agric Food Chem* 43:1659-68.
- Scheiberle P, Hofman T. 1997. Evaluation of the character impact odorants in fresh strawberry juice by quantitative measurements and sensory studies on model mixtures. *J Agric Food Chem* 45:227-32.
- Siverstein HK, Risvik E. 1994. A study of sample and assessor variation: a multivariate study of wine profiles. *J Sens Stud* 9:293-312.
- Stampanoni CR. 1994. The use of standardized flavor languages and quantitative flavor profiling technique for flavored dairy products. *J Sens Stud* 9:383-400.
- Stone H. 1992. Quantitative descriptive analysis. In: Hootman RC, editor. *ASTM Manual Series MNL 13 Manual on Descriptive Analysis Testing for Sensory Evaluation*. West Conshohocken, Pa.: Am. Soc. Testing and Materials. p 15-21.
- Suriyaphan O, Drake MA, Cadwallader KR. 1999. Identification of volatile off-flavors in reduced-fat Cheddar cheeses containing lecithin. *Lebensm Wiss U Technol* 32(5):250-4.
- Suriyaphan O, Drake MA, Chen XQ, Cadwallader KR. 2001. Characteristic aroma components of British Farmhouse Cheddar cheese. *J Agric Food Chem* 49:1382-7.
- Taylor AJ. 1996. Volatile flavor release from foods during eating. *Crit Rev Food Sci Nutr* 36:765-84.
- Van Ruth SM, O'Connor CH. 2001. Evaluation of 3 gas chromatography-olfactometry methods: comparison of odour intensity-concentration relationships of 8 volatile compounds with sensory headspace data. *Food Chem* 74:341-7.
- Van Ruth SM, Grossmann I, Geary M, Delahunty CM. 2001. Interactions between artificial saliva and 20 aroma compounds in water and oil model systems. *J Agric Food Chem* 49:2409-13.
- Virgili R, Parolari G, Bolzoni L, Barbieri G, Mangia A, Careri M, Spagnoli S, Panari G, Zannoni M. 1994. Sensory chemical relationships in Parmigiano-Reggiano Cheese. *Lebensm Wiss Technol* 27:491-5.
- Watson MP, McEwan JA. 1995. Sensory changes in liquid milk during storage and the effect on consumer acceptance. *J Soc Dairy Technol* 48:1-8.
- Wu YF, Baek HH, Gerard PD, Cadwallader KR. 2000. Development of a meat-like process flavoring from soybean based enzyme hydrolyzed vegetable protein (E-HVP). *J Food Sci* 65:1220-7.
- Yau NJN, Liu TT. 1999. Instrumental and sensory analysis of volatile aroma of cooked rice. *J Sens Stud* 14:209-33.
- Young H, Gilbert JM, Murray SH, Ball RD. 1996. Causal effects of aroma compounds on Royal Gala apple flavours. *J Sci Food Agric* 71:329-36.
- Zook K, Wessman C. 1977. The selection and use of judges for descriptive panels. *Food Technol* 31:56-61.

MS 20010013 Submitted 1/8/02, Accepted 4/9/02, Received 4/10/02

Author Drake gratefully acknowledges the support of Dairy Management, Inc., California Dairy Research Foundation, and North Carolina State Univ. This manuscript is paper nr 02-52 in the journal series of the Dept. of Food Science, North Carolina State Univ., Raleigh, N.C. The use of trade names in this publication does not imply endorsement by the North Carolina Agricultural Research Service nor criticism of similar ones not mentioned.

Author Drake is with the Dept. Food Science, North Carolina State Univ., Raleigh, N.C., U.S.A. Author Civile is with Sensory Spectrum, Inc., Chatham, N.J., U.S.A. Direct inquiries to author Drake (E-mail: mdrake@unity.ncsu.edu).